

The maximum momentum of particles accelerated at cosmic ray modified shocks

Pasquale Blasi^{1*}, Elena Amato^{1†} and Damiano Caprioli^{2‡}

¹*INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125, Firenze, Italy*

²*Scuola Normale Superiore, Pisa, Italy*

Accepted —. Received —

ABSTRACT

Particle acceleration at non-relativistic shocks can be very efficient, leading to the appearance of non-linear effects due to the dynamical reaction of the accelerated particles on the shock structure and to the non-linear amplification of the magnetic field in the shock vicinity. The value of the maximum momentum p_{max} in these circumstances cannot be estimated using the classical results obtained within the framework of test particle approaches. We provide here the first attempt at estimating p_{max} in the cosmic ray modified regime, taking into account the non-linear effects mentioned above.

Key words: acceleration of particles - shock waves

1 INTRODUCTION

Since the beginning of the investigations on the mechanism of particle acceleration at shock fronts, estimating the maximum energy of the accelerated particles has been one of the main goals, in the perspective of assessing the ability of the mechanism to explain the observed energies in cosmic rays. This issue has played a crucial role especially in the context of the so-called supernova paradigm for the origin of galactic cosmic rays. In fact, it was understood very soon, in the case of supernova remnants' shocks, that acceleration would not be efficient enough and provide sufficiently high energies unless magnetic scattering

* E-mail: blasi@arcetri.astro.it

† E-mail: amato@arcetri.astro.it

‡ E-mail: caprioli@sns.it

could be self-generated by the particles themselves through their motion in the upstream plasma. The mechanism through which this generation of waves could occur was identified in the streaming instability (Bell (1978a)) and widely discussed later by Lagage & Cesarsky (1983a,b). In these latter papers, in particular, the maximum energy was estimated for shocks in supernova remnants, already believed to be candidate sources of galactic cosmic rays. The result of this investigation was however that the maximum energy, even in the presence of self-generated scattering agents, is of the order of $\sim 10^3 - 10^4$ GeV, quite lower than the energy at which the knee is observed. Today we know that this energy is also lower than the maximum energy observed in the proton component by the KASCADE experiment (see Hörandel (2005) for a comprehensive review of experimental results in the knee region) that seems to be as high as $\sim 10^7$ GeV (see Blasi (2005) for a recent review).

More recently the problem has been complicated even more by the understanding that particle acceleration can be rather efficient and break the *test particle* regime that old calculations were based upon. This important result was obtained independently within completely different approaches, two-fluid models (Drury & Völk (1980, 1981)), kinetic models (Malkov (1997); Malkov, Diamond & Völk (2000); Blasi (2002, 2004); Amato & Blasi (2005, 2006)) and numerical approaches, using both Monte Carlo and other simulation procedures (Jones & Ellison (1991); Bell (1987); Ellison, Möbius & Paschmann (1990); Ellison, Baring & Jones (1995, 1996); Kang & Jones (1997, 2005); Kang, Jones & Gieseler (2002); Vladimirov, Ellison & Bykov (2006)). The non-linear dynamical reaction of the accelerated particles introduces a positive reaction that results in a flattening of the high energy spectra, which are no longer power laws, thereby enhancing the role of particles at the highest energies. Moreover a gradient in the velocity of the upstream fluid is produced by the pressure of the accelerated particles. The value of p_{max} is important to determine the spectrum, and the spectrum determines the amplification of the magnetic field which in turn determines p_{max} . The formalism of Lagage & Cesarsky (1983a,b) is not appropriate to describe this type of situation.

In this paper we describe the first calculation of the maximum momentum for cosmic ray modified shocks, including all the relevant non-linear effects. The description of the acceleration process, with the dynamical reaction of cosmic rays and the magnetic field amplification taken into account, is based on the recent papers by Amato & Blasi (2005, 2006).

The paper is organized as follows: in §2 we describe the method that allows us to compute the acceleration time as a function of the particle momentum; in §3 we describe the

procedure used to determine p_{max} in the case of modified shocks and discuss the underlying assumptions; in §4 we present the results we find for the maximum achievable momentum in supernova remnant shocks; finally a critical discussion of the method adopted and a summary of the main results is provided in § 5.

2 ACCELERATION TIME FOR MODIFIED SHOCKS

The full description of the calculations that allow us to compute the spectrum of the accelerated particles, the dynamics of the upstream and downstream plasmas and the strength of the amplified magnetic field due to cosmic rays' streaming instability in the stationary regime can be found in the papers by Amato & Blasi (2005, 2006).

In this section we present the main achievement of our work, which consists of a derivation of the acceleration time $t_{acc}(p)$ (as a function of the momentum p) for particles accelerated at a shock with an arbitrary level of shock modification. The expression we derive is a generalization of the well known formulae found in the context of quasi-linear theory (Lagage & Cesarsky (1983a,b)) and reduces to them when the non-linear shock modification is negligible.

In general, the transport of the accelerated particles is described by the time dependent diffusion-convection equation:

$$\frac{\partial f}{\partial t} + u \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} \left[D \frac{\partial f}{\partial x} \right] + \frac{1}{3} \frac{du}{dx} p \frac{\partial f}{\partial p} + Q(x, p), \quad (1)$$

where $f(t, x, p)$ is the distribution function of the accelerated particles, $D(t, x, p)$ is the spatial diffusion coefficient and Q is the injection term, taken in the form $Q(x, p) = Q_0(p)\delta(x)$, with the assumption that injection takes place only at the shock surface on the downstream side (although this assumption is not strictly needed). The x -axis goes from upstream infinity ($x = -\infty$) to downstream infinity ($x = +\infty$) and the shock is at $x = 0$. We will use the index 1 (2) to refer to the region immediately upstream (downstream) of the shock.

The problem of determining the time scale associated with the change of the distribution function around the maximum momentum is impossible to face analytically in all its complexity, as implied by the non-linearity of the system: in general, f , u and D all depend on time, and change on comparable time scales. Moreover, f depends in a complicated way on all other quantities (Amato & Blasi (2005, 2006)).

Let us assume that we know an approximation of the solution f in the form of a solution of the stationary problem for a given value of the maximum momentum p_{max} , as can be

calculated following Amato & Blasi (2005, 2006). This value of p_{max} can be considered as a *guess* value to start with. In this case the velocity profile $u(x)$ can be considered as determined in first instance from the guess function f as derived by the assumption of quasi-stationarity, using the calculations of Amato & Blasi (2005, 2006). Within this approximate framework, we can introduce the Laplace transform of the distribution function with respect to time:

$$g(s, x, p) = \int_0^\infty dt e^{-st} f(t, x, p), \quad (2)$$

so that the transport equation becomes:

$$sg + u \frac{\partial g}{\partial x} = \frac{\partial}{\partial x} \left[D \frac{\partial g}{\partial x} \right] + \frac{1}{3} \frac{du}{dx} p \frac{\partial g}{\partial p} + \frac{1}{s} Q(x, p), \quad (3)$$

Integration of Eq. 3 in the vicinity of the shock (namely between $x = 0^-$ and $x = 0^+$) leads to the following boundary condition at the shock:

$$\left[D \frac{\partial g}{\partial x} \right]_2 - \left[D \frac{\partial g}{\partial x} \right]_1 + \frac{1}{3} (u_2 - u_1) p \frac{\partial g_0(s, p)}{\partial p} + \frac{1}{s} Q_0(p) = 0, \quad (4)$$

where $u_1 = u(0^+)$, $g_0(s, p) = g(s, x = 0, p)$ and we have used the continuity of the distribution function (at any time) at the shock.

We can find another condition on the distribution function by integrating the transport equation (Eq. 3) in the upstream section from $x = -\infty$ to $x = 0^-$, so as to obtain:

$$sI(s, p) + u_1 g_0(s, p) - \int_{-\infty}^{0^-} dx g(s, x, p) \frac{du(x)}{dx} = \left[D \frac{\partial g}{\partial x} \right]_1 + \frac{1}{3} p \frac{\partial}{\partial p} \int_{-\infty}^{0^-} dx g(s, x, p) \frac{du(x)}{dx}, \quad (5)$$

where $I(s, p) = \int_{-\infty}^{0^-} dx g(s, x, p)$. Following Blasi (2002) we introduce the quantity

$$u_p(s, p) = u_1 - \frac{1}{g_0} \int_{-\infty}^{0^-} dx g(s, x, p) \frac{du}{dx}, \quad (6)$$

which allows us to rewrite Eq. 5 as

$$\left[D \frac{\partial g}{\partial x} \right]_1 = sI(s, p) + g_0 u_p \left(1 + \frac{1}{3} \frac{d \ln u_p}{d \ln p} \right) - \frac{1}{3} p \frac{\partial g_0}{\partial p} (u_1 - u_p). \quad (7)$$

Clearly in this equation u_p depends on time through the functions f and u , which again reflects the non-linearity of the underlying equation.

The transport equation in the downstream fluid can be considerably simplified if we assume that the fluid velocity is independent of location $u(x) = u_2$ and that the diffusion coefficient is also independent of x . In this case, for any $x \neq 0$ Eq. 1 becomes:

$$\frac{\partial f}{\partial t} + u_2 \frac{\partial f}{\partial x} = D_2 \frac{\partial^2 f}{\partial x^2}. \quad (8)$$

If we use again the Laplace transform of the distribution function with respect to time we have

$$sg + u_2 \frac{\partial g}{\partial x} = D_2 \frac{\partial^2 g}{\partial x^2}, \quad (9)$$

whose solution has the form $g(s, x, p) = g_0(p)e^{-\beta x}$ (in the downstream region $x > 0$) with β easily found by substitution, so that:

$$g(s, x, p) = g_0(p) \exp \left\{ \frac{u_2 x}{2D_2} \left[1 - \left(1 + \frac{4sD_2}{u_2^2} \right)^{1/2} \right] \right\}. \quad (10)$$

This implies that

$$\left[D \frac{\partial g}{\partial x} \right]_2 = \frac{1}{2} g_0(p) u_2 \sigma_2, \quad \sigma_2 = 1 - \left(1 + \frac{4sD_2}{u_2^2} \right)^{1/2}. \quad (11)$$

At this point, substituting Eqs. 7-11 into the jump condition Eq. 4 we obtain the following equation:

$$g_0 \left\{ \frac{1}{2} u_2 \sigma_2 - u_p \left(1 + \frac{1}{3} \frac{d \ln u_p}{d \ln p} \right) \right\} + \frac{1}{3} (u_2 - u_p) p \frac{dg_0}{dp} - sI + \frac{1}{s} Q_0 = 0. \quad (12)$$

As already proposed by Malkov (1997) and Amato & Blasi (2005), it is useful to rewrite the transport equation in an implicit form by integrating it between $x = -\infty$ and a generic point x upstream. We repeat this procedure here on the Laplace-transformed transport equation (Eq. 3). This leads to the following implicit expression for g :

$$g(s, x, p) = \exp \left\{ \int_0^x dx' \frac{u(x')}{D(x', p)} \right\} \left\{ g_0(s, p) + \int_0^x dx' \frac{1}{D(x', p)} \exp \left[- \int_0^{x'} dx'' \frac{u(x'')}{D(x'', p)} \right] \times \right. \\ \left. \left\{ - \int_{-\infty}^{x'} dx'' g(s, x'', p) \frac{du}{dx''} \left(1 + \frac{1}{3} \frac{d \ln g(s, x'', p)}{d \ln p} \right) + s \int_{-\infty}^{x'} dx'' g(s, x'', p) \right\} \right\}. \quad (13)$$

As already discussed by Malkov (1997) and Amato & Blasi (2005), one can approximate this complex implicit solution as

$$g(s, x, p) = g_0(s, p) \exp \left\{ \frac{q(p)}{3} \int_0^x dx' \frac{u(x')}{D(x', p)} \right\} + \\ + s \exp \left[\int_0^x dx' \frac{u(x')}{D(x', p)} \right] \int_0^x dx' \frac{1}{D(x', p)} \int_{-\infty}^{x'} dx'' g(x'') \quad (14)$$

where $q(p)$ is now the slope of the distribution function at the shock. This expression is obtained by expanding the previous expression for small values of $\int_0^x dx' \frac{u(x')}{D(x', p)}$. In Amato & Blasi (2005) it was shown *a posteriori* that this is a good approximation of the spectral distribution of the particles as a function of the spatial coordinate.

The agreement was found to be worse when the shock is quasi-linear, namely for small modifications. The reason for this is in the fact that Eq. 14 satisfies the jump condition in the stationary regime ($s \rightarrow 0$) only when $u_2 \rightarrow 0$. A better approximation for g is in fact of

the form,

$$g(s, x, p) = g_0(s, p) \exp \left\{ \frac{q(p)}{3} \left(1 - \frac{u_2}{u_1} \right) \int_0^x dx' \frac{u(x')}{D(x', p)} \right\} + \\ + s \exp \left[\int_0^x dx' \frac{u(x')}{D(x', p)} \right] \int_0^x dx' \frac{1}{D(x', p)} \int_{-\infty}^{x'} dx'' g(x'') \quad (15)$$

which satisfies the jump condition exactly. Note that Eq. 15 differs from Eq. 14 only in that at given momentum the particle distribution is slightly more extended spatially. In the following we adopt Eq. 15 as an approximation of the spatial distribution of the accelerated particles. In evaluating the term $sI(s, p)$ in Eq. 12 we retain only the first term of g in Eq. 15, since in the end we will be interested in the limit $s \rightarrow 0$ (large times). This means that Eq. 12 becomes:

$$g_0 \left\{ \frac{1}{2} u_2 \sigma_2 - s \Lambda(p) - u_p \left(1 + \frac{1}{3} \frac{d \ln u_p}{d \ln p} \right) \right\} + \frac{1}{3} (u_2 - u_p) p \frac{dg_0}{dp} + \frac{1}{s} Q_0 = 0, \quad (16)$$

where

$$\Lambda(p) = \int_{-\infty}^0 dx \exp \left\{ \frac{q(p)}{3} \left(1 - \frac{u_2}{u_1} \right) \int_0^x dx' \frac{u(x')}{D(x')} \right\}. \quad (17)$$

The solution of this equation is easily found to be:

$$g_0(s, p) = \frac{1}{s} \frac{\eta n_{gas,1}}{4\pi p_{inj}^3} \frac{3R_{sub}}{R_{tot} U_p(p) - 1} \exp \left\{ - \int_{p_{inj}}^p \frac{dp'}{p'} \frac{3U_p(p') R_{tot}}{R_{tot} U_p(p') - 1} \right\} \times \\ \times \exp \left\{ \int_{p_{inj}}^p \frac{dp'}{p'} \frac{3}{2} \frac{\sigma_2}{R_{tot} U_p(p') - 1} \right\} \times \exp \left\{ - \int_{p_{inj}}^p \frac{dp'}{p'} \frac{3s \Lambda(p')}{R_{tot} U_p(p') - 1} \right\}, \quad (18)$$

or, in a more compact form,

$$g_0(s, p) = \frac{1}{s} K(p) \exp [h(p, s)], \quad (19)$$

where

$$K(p) = \frac{\eta n_{gas,1}}{4\pi p_{inj}^3} \frac{3R_{sub}}{R_{tot} U_p(p) - 1} \exp \left\{ - \int_{p_{inj}}^p \frac{dp'}{p'} \frac{3U_p(p') R_{tot}}{R_{tot} U_p(p') - 1} \right\} \quad (20)$$

and

$$h(p, s) = \int_{p_{inj}}^p \frac{dp'}{p'} \left[\frac{3}{2} \frac{\sigma_2}{R_{tot} U_p(p') - 1} - \frac{3R_{tot}}{u_0} \frac{s \Lambda(p')}{R_{tot} U_p(p') - 1} \right], \quad (21)$$

with $U_p(p) = u_p/u_0$ and $u_0 = u(-\infty)$.

At this point we follow the elegant procedure discussed by Drury (1983). We need to calculate the inverse Laplace transform

$$f(t, x, p) = \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} ds \frac{e^{ts}}{s} K(p) e^{h(p,s)}.$$

We introduce the function

$$\phi(p, t) = \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} ds e^{ts} e^{h(p,s)}. \quad (22)$$

One can easily show that the inverse Laplace transform of the function

$$\tilde{f}(p, t) = K(p) \int_0^t dt' \phi(p, t') \quad (23)$$

is exactly the function g_0 as written in Eq. 19. It follows that the solution of our problem is in fact $f = \tilde{f}$. In the approximate framework in which we are, the function ϕ , as in the linear case discussed by Drury (1983), represents the probability that a particle starting with energy p_{inj} has reached the momentum p in a time t (ϕ is normalized to unity). The mean time for the acceleration to momentum p is therefore $\langle t(p) \rangle = \int_0^\infty dt t \phi(p, t)$. One can use the following obvious property of the function ϕ

$$\int_0^\infty dt \phi(p, t) e^{-ts} = \exp[h(p, s)], \quad (24)$$

to determine $\langle t \rangle$, by deriving Eq. 24 with respect to s and calculating it at $s = 0$. From this follows that:

$$\langle t \rangle = - \left[\frac{\partial h}{\partial s} \right]_{s=0} = \frac{3R_{tot}}{u_0^2} \int_{p_{inj}}^p \frac{dp'}{p'} \left\{ \frac{R_{tot}D_2(p')}{R_{tot}U_p(p') - 1} + \frac{u_0\Lambda(p')}{R_{tot}U_p(p') - 1} \right\}. \quad (25)$$

For an unmodified shock, with spatially constant fluid velocities upstream and downstream, one has that $q(p) = 3u_1/(u_1 - u_2)$ and the previous formula easily reduces to the well known:

$$\langle t \rangle = \frac{3}{u_1 - u_2} \left[\frac{D_2}{u_2} + \frac{D_1}{u_1} \right], \quad (26)$$

if the diffusion coefficient is in the form of Bohm diffusion. For an arbitrary form of the diffusion coefficient the expression in Eq. 26 is correct within a factor of order unity.

3 CALCULATION OF P_{MAX}

In general the value of p_{max} is determined by the competition between the acceleration time and the shortest among the finite age of the source, the time scale for energy losses, and the time scale for escape of particles with momentum p_{max} from upstream infinity.

Here we concentrate on the self-consistent calculation of the maximum momentum in those cases when it is determined by the finite age of the source (this is expected to be the case for supernova remnants). In such a context each value of p_{max} therefore reflects a different age τ of the system. In this sense, our previous work (Amato & Blasi (2005, 2006)) was based on the implicit assumption that a quasi-stationary regime could be reached on time scales shorter than τ , but no specific recipe for the calculation of p_{max} was adopted. In other work on non-linear diffusive particle acceleration at shock waves, p_{max} was either fixed or determined according with a simple recipe based on test-particle theory.

Our determination of the maximum momentum proceeds through 2 different steps: 1) we first determine the acceleration time $t_{acc}(p)$ (Eq. 25) for a given value of $p_{max}^{(k)}$ (at the first step this is just a guess value $p_{max}^{(0)}$); 2) we compare $t_{acc}(p_{max}^{(k)})$ with the age of the system τ and determine an updated value of p_{max} , say $p_{max}^{(k+1)}$. The procedure is iterated until $t_{acc}(p_{max}) = \tau$, where p_{max} is now the actual maximum momentum for our problem. It is crucial to keep in mind that due to the non-linearity of the system, changing p_{max} causes the all system to change (velocity profile, spectrum of accelerated particles, efficiency of particle acceleration and also the diffusion coefficient, if the scattering is due to self-generated waves).

The results obtained by applying this procedure are illustrated in the next section.

4 RESULTS

Following the method outlined in § 2 we computed the acceleration time as a function of the particle momentum p for a system with $\tau = 1000 \text{ yr}$, under different assumptions on the shock Mach number at infinity M_0 and on the diffusion properties of the system. The results of these calculations are summarized in Fig. 1. We assumed a velocity of the shock wave of $5 \times 10^3 \text{ km/s}$ and considered values of M_0 ranging from 5 (unmodified shocks) to 200 (already in the asymptotic regime for strongly modified shocks). The injection momentum was taken to be ξp_{therm} with p_{therm} corresponding to the maxwellian temperature T_2 and $\xi = 3.5$. We also related the injection efficiency η to the compression ratio at the subshock through the recipe proposed by Blasi, Gabici & Vannoni (2005) and already adopted by Amato & Blasi (2005, 2006). In the top panel of Fig. 1 we plot the acceleration time up to a momentum p for $D(x, p) = D_{sg}$, namely with self-generated scattering, with $D_{sg} = (4/3\pi)vr_L/\mathcal{F}$ the result of quasi-linear theory for scattering induced by self-generated Alfvén waves. Here v and r_L are the particle velocity and Larmor radius (in the background field B_0) respectively, and $\mathcal{F}$ is the dimensionless wave energy density per unit logarithmic band-width (see e.g. Amato & Blasi (2006) for details). In the bottom panel of Fig. 1 we show the acceleration time for $D(x, p) = D_{Bohm}$, with D_{Bohm} referring to Bohm-like diffusion in the amplified magnetic field δB found by integrating $B_0^2 \mathcal{F}$ in k -space: $D_{Bohm} = (1/3)vr_L(B_0/\delta B)$. In all of our calculations the background magnetic field was taken to be $B_0 = 10\mu G$.

The maximum momentum obtained from our iterative procedure is seen as the intersection with the horizontal line $\tau = 1000$ years, assumed to be the age of the system.

Each of the curves in Fig. 1 was obtained by following the procedure outlined in § 3. The




Figure 1. We plot the time (in years) it takes for the system to accelerate particles up to a momentum p (in units of mc) under different conditions. The different panels refer to different choices of the diffusion coefficient. In the top panel the diffusion coefficient is the self-generated one, while in the bottom panel we assume Bohm diffusion in the amplified magnetic field (see text for details). The different line types refer to different values of the shock Mach number: $M_0 = 5$ for the solid line, $M_0 = 20$ for the dotted line, $M_0 = 100$ dashed, $M_0 = 200$ for the dot-dashed line.

first thing to notice by comparing the two different panels in Fig. 1 is that the maximum achievable momentum, for a given Mach number, is roughly the same (within a factor of a few (2-4) at most) for the two diffusion models adopted. This observation is very important given the fact that the diffusion coefficient is basically unknown and not easily obtained from observations.

Still the comparison between the two panels allows to highlight what the main effect of the different diffusion models is: in the case of strongly modified shocks the spectra become very flat at high energies as discussed in detail elsewhere (e.g. Amato & Blasi (2006)) and most of the power of the Alfvén waves excited by the streaming cosmic rays is deposited in long wavelength waves (the ones that are resonant with high energy particles), with a relatively smaller fraction going into waves that are able to efficiently scatter the low energy particles. This is the reason for the behaviour of $t_{acc}(p)$ observed in the upper panel, which may seem surprising at first sight for the flatness of the curves referring to $M_0 = 100 - 200$.

By comparing the two panels, it is clear that things change drastically, with the dependence of t_{acc} on p becoming much stronger, if the energy of the waves is redistributed uniformly between different wavelengths, which is what one is effectively assuming by adopting a Bohm-like diffusion coefficient in the amplified magnetic field.

Finally, let us discuss the behaviour of the maximum achievable momentum as a function of the Mach number. This is qualitatively the same for both models of diffusion: p_{max} initially increases quickly with the Mach number as a result of field amplification in modified shocks and then decreases somewhat, although slowly, when M_0 is further increased. This latter fact is clearly related to the more strongly modified velocity profile that leads to lower and lower values of the flow velocity in the vicinity of the shock and as a consequence increasingly longer residence times upstream. The dependence of the maximum achievable momentum on the shock Mach number is illustrated more clearly in Fig. 2, where we plot p_{max} as a function of M_0 for four different models of diffusion. The plot refers again to a system with $\tau = 10^3$ yr and all other values of the relevant parameters are the same as above, including the background magnetic field strength and the recipe for p_{inj} .

The solid curve refers to $D(x, p) = D_{sg}(x, p)$ as described above, the dashed curve is for $D(x, p) = D_{Bohm}(x, p)$, the dotted curve is for $D(x, p) = D_{Bohm}(0, p)$ (namely the diffusion coefficient equals the Bohm diffusion calculated in the amplified magnetic field at the shock surface, without spatial variation) and finally the dot-dashed curve refers to Bohm diffusion in the background magnetic field. The first aspect that clearly appears from the figure is the enhancement of p_{max} that arises from magnetic field amplification. For high Mach number shocks the magnetic field strength at the shock is amplified by a factor of order 25 and this implies an enhancement by a factor 10-30 (depending on the diffusion model adopted) of the maximum achievable energy (comparison of the solid, dashed or dotted curves with the dot-dashed one). On the other hand, p_{max} does not strongly depend on the detailed diffusion model adopted, in terms of the turbulence spectrum, with the difference between Bohm diffusion and resonant scattering with self-generated Alfvén turbulence (comparison between the solid and dashed curves) being a factor of 2 at most. A few remarks are worth being made. Note that the values of the parameters adopted throughout our calculations are reminiscent of those appropriate to the case of young supernova remnants (SNRs). This is because we wished to address here the issue of the maximum energy that can be achieved in this class of sources, as candidate sources of galactic cosmic rays. Moreover there seems to be now solid evidence for amplified magnetic fields in SNRs (Warren (2005)), so that there is




Figure 2. We plot the maximum momentum the particles can achieve within a time $\tau = 10^3$ yr as a function of the Mach number of the shock. The different curves refer to different choices of the diffusion coefficient: the solid line is for self-generated diffusion; the dashed line is for Bohm diffusion in the self-amplified magnetic field; the dotted line is for Bohm diffusion in a magnetic field that is spatially constant in strength and equal to $\delta B(0)$, the value reached at the shock location by the cosmic ray streaming induced field; finally the dot-dashed line is for Bohm diffusion in the background magnetic field.

observational support for the idea that cosmic rays might be efficiently accelerated at SNRs' shocks. The shock modification is a natural consequence of the effectiveness of these shocks as cosmic ray accelerators.

The maximum energy that the accelerated particles can achieve in the context of modified shocks is always less than one would naively guess by using a Lagage & Cesarsky (1983a) type of estimate with the upstream velocity set equal to u_0 and the magnetic field evaluated at the shock surface. This is due to the combination of two effects. On one side, the net effect of the shock modification is to slow down the upstream fluid (precursor), so that the effective velocity *felt* by the particles of given momentum is always lower than u_0 , thereby making the acceleration process slower. This is illustrated effectively through the dotted line in Fig. 2, where the diffusion coefficient is taken as Bohm-like calculated with the magnetic field at the shock, assumed constant in the precursor. In this case the value of p_{max} is always lower than $\sim 10^6$ GeV/c for Mach number $M_0 > 100$, while in the absence of shock modification

one would get $p_{max} \sim 6 \times 10^6$ GeV/c. This factor > 5 is solely due to the existence of a precursor. There is however a second (less important) effect that causes a further decrease of p_{max} : the spatial distribution of the accelerated particles in the precursor also determines a spatial profile of the strength of the magnetic field associated with the self-generated Alfvén waves, so that the effective diffusion coefficient is also larger than $D_{Bohm}(0, p)$. This effect is visible when comparing the dashed line with the dotted line in Fig. 2. The combination of the two effects leads to an overall decrease of p_{max} by a factor ~ 10 compared with the naive expectation *a la* Lagage & Cesarsky (1983a,b), more evident for higher values of the Mach number.

5 A CRITICAL DISCUSSION OF THE METHOD AND SUMMARY OF THE MAIN RESULTS

In this section we summarize the reasons for the need of the type of calculations presented here and we discuss some shortcomings of the proposed mathematical approach.

To our knowledge this represents the first attempt to determine the maximum momentum of the particles accelerated at a cosmic ray modified shock, where the nonlinear dynamical reaction of the accelerated particles changes the shocked fluid, the particles self-generate their own scattering centers, and the modified system in turn determines the spectral shape and energy content of the accelerated particles' distribution. The creation of a shock precursor in the upstream region and the amplification of the background magnetic field through streaming instability are the main reasons to expect that the maximum achievable momentum may be appreciably different from the one calculated in linear theory for the same value of the shock Mach number. The precursor, due to the gradual slowing down of the upstream plasma would by itself be expected to slightly decrease the maximum momentum: this effect can in fact be seen in Fig. 2, where the dash-dotted line shows the maximum momentum as a function of the Mach number when the background field is not amplified. The reduced velocity in the upstream plasma slows down the process of return to the shock for the particles in the upstream section of the fluid. In the presence of magnetic field amplification the maximum momentum is expected to increase. This trend continues up to Mach numbers for which the pressure in the form of accelerated particles cannot increase any more (saturation), so that the self-generated magnetic field also saturates. On the other hand the compression factor (shock modification) keeps increasing, so that the corresponding p_{max} in

Fig. 2 shows a slightly decreasing trend (roughly the same that one obtains in the presence of shock modification but without field amplification, dash-dotted line). One should be careful not to mistake this trend for a temporal trend. The behaviour of the maximum momentum is to be interpreted only as a trend as a function of the fluid Mach number at upstream infinity. In a more realistic situation the Mach number can also change with time, which might introduce additional effects, not taken into account here. On the other hand, if the conclusions of Lagage & Cesarsky (1983b) remain valid for this more complex scenario, one should expect that most of the acceleration of cosmic rays occurs between the end of the free expansion phase and the beginning of the Sedov phase. This is the phase we concentrated our attention upon, so that the assumption of temporally constant Mach number adopted here is justified.

The most important equation derived in this work is Eq. 25 for the acceleration time in the presence of shock modification. It is important to realize that although the calculations shown here have been carried out by adopting the theory of Amato & Blasi (2005, 2006) for the evaluation of the modification and field amplification, Eq. 25 can be applied in a very general way. In fact all the quantities involved in the evaluation of Eq. 25 can be determined using very different approaches, for instance Monte Carlo simulations, while Eq. 25 can be adopted to estimate the *instantaneous* value of p_{max} . Moreover, since Eq. 25 directly contains the diffusion coefficients and the dynamical quantities (e.g. fluid velocity), it can be applied independently of the specific mechanism of wave production. If the non-resonant wave generation mechanism recently proposed by Bell (2004) were to be responsible for the wave production and particle scattering, this would change the mathematical form of the diffusion coefficient, and possibly the dynamical reaction of the plasma (if the amplified field reaches a dynamically important value), but Eq. 25 could still be applied unchanged.

The most relevant shortcoming of the calculation illustrated here is in the fact that we used the assumption of quasi-stationarity in the upstream fluid (not downstream). This assumption is fundamental to allow us to apply the method of Laplace transform, which can be used only for linear differential equations. At the same time quasi-stationarity is also required in order to use our previous calculations (Amato & Blasi (2005, 2006)) for the shock modification and field amplification. These are actually two different aspects of the same problem of this approach, which however cannot be avoided if an analytical attempt has to be made to calculate the maximum momentum in these complex situations.

It should however be stressed that the shortcomings just discussed also apply to the

approach of Lagage & Cesarsky (1983a,b): their calculations are only valid for linear shocks ($P_{CR} \ll \rho_0 u_0^2$) so that the velocity and density profiles at the shock are spatially constant and given. On the other hand the diffusion coefficient, which is self-generated in their approach, is time-dependent because the particle distribution function is time-dependent. Despite this, the authors fix the diffusion coefficient (and the distribution function of the accelerated particles) and proceed by applying the method of the Laplace transform. In this sense our approach should be considered as the extension of the work of Lagage & Cesarsky (1983a,b) to the case in which the dynamical reaction of the accelerated particles is not neglected. The common finding of several independent approaches to particle acceleration at shocks is that shock modification manifests itself in a prominent way, as a consequence of efficient acceleration, therefore our extension seems particularly important at this time.

We conclude this section by providing a short, itemized summary of our main scientific results:

1) the maximum momentum of accelerated particles can be as high as $\sim 10^6$ GeV/c for protons, when all effects are taken into account. The approach of Lagage & Cesarsky (1983a,b) would give $p_{max} \sim 6 \times 10^6$ GeV/c when the magnetic field is assumed spatially constant to its maximum value at the shock surface. The same approach for $\delta B/B_0 \approx 1$ would reproduce the well known value $p_{max} = 10^5$ GeV/c for $B_0 = 10\mu G$ and $u_0 = 5000\text{km/s}$.

2) the net effect of the shock modification is to create a precursor in both the fluid velocity and magnetic field strength. The former decreases while moving towards the shock while the latter increases. Both phenomena contribute to lowering p_{max} with respect to the value $\sim 6 \times 10^6$ GeV/c.

3) Provided the magnetic field is amplified by the streaming instability of cosmic rays, the value of p_{max} is weakly dependent upon the exact spectrum of the magnetic turbulence.

4) The expression provided in Eq. 25 is the only expression presently available that allows one to calculate the acceleration time for cosmic ray modified shock waves. This expression is very general and can be applied independently of the specific treatment of the shock modification and reduces to the well known formula for weakly modified shocks.

REFERENCES

- Amato, E., and Blasi, P., 2005, MNRAS Lett., 364, 76
 Amato, E., and Blasi, P., 2006, MNRAS, in press

- Bell, A.R., 1978a, MNRAS, 182, 147
- Bell, A.R., 1987, MNRAS, 225, 615
- Bell, A.R., 2004, MNRAS 353, 550.
- Blasi, P., 2002, Astropart. Phys. 16, 429
- Blasi, P., 2004, Astropart. Phys. 21, 45
- Blasi, P., 2005, Modern Phys. Lett. A, 20, 1
- Blasi, P., Gabici, S., and Vannoni, G., 2005, MNRAS, 361, 907
- Drury, L.O'C and Völk, H.J., 1981, in Proc. IAU Symp. 94 on "Origin of Cosmic Rays", Setti G., Spada G., Wolfendale A. W., eds, Reidel Publishing, Dordrecht, p.363
- Drury, L.O'C and Völk, H.J., 1981, ApJ, 248, 344
- Drury, L .O'C., 1983, Rep. Prog. Phys. 46, 973
- Ellison, D.C., Möbius, E., and Paschmann, G., 1990, ApJ, 352, 376
- Ellison, D.C., Baring, M.G., and Jones, F.C., 1995, ApJ, 453, 873
- Ellison, D.C., Baring, M.G., and Jones, F.C., 1996, ApJ, 473, 1029
- Hörandel, J.R., *A review of experimental results at the knee*, in the "Workshop on Physics of the End of the Galactic Cosmic Ray Spectrum", Aspen, April 25 - 29, 2005 (preprint astro-ph/0508014).
- Jones, F.C. and Ellison, D.C., 1991, Space Sci. Rev. 58, 259
- Kang, H., and Jones, T.W., 1997, ApJ, 476, 875
- Kang, H., Jones, T.W., and Gieseler, U.D.J., 2002, ApJ 579, 337
- Kang, H., and Jones, T.W., 2005, ApJ, 620, 44
- Lagage, P.O., and Cesarsky, C.J., 1983, A&A, 118, 223
- Lagage, P.O., and Cesarsky, C.J., 1983, A&A, 125, 249
- Malkov, M.A., 1997, ApJ, 485, 638
- Malkov, M.A., Diamond P.H., and Völk, H.J., 2000, ApJL, 533, 171
- Vladimirov, A., Ellison, D.C., and Bykov, A., 2006, ApJ, 652, 1246
- Warren, J.S., et al., 2005, ApJ, 634, 376.